

# From Inflation to Dark Energy

Robert Brout

*Faculté des Sciences, Université Libre de Bruxelles, Belgium  
and Perimeter Institute for Theoretical Physics  
Waterloo, Ontario, Canada*

It is proposed that after the macroscopic fluctuation of energy density that is responsible for inflation dies away, a class of microscopic fluctuations, always present, survives to give the present day dark energy. This latter is simply a reinterpretation of the causet mechanism of Ahmed, Dodelson, Green and Sorkin, wherein the emergence of space is dropped but only energy considerations are maintained. At postinflation times, energy is exchanged between the "cisplanckian" cosmos and an unknown foam-like transplanckian reservoir. Whereas during inflation, the energy flows only from the latter to the former after inflation it fluctuates in sign thereby accounting for the tiny effective cosmological constant that seems to account for dark energy.

It has become apparent in recent years that the most important component of energy that drives the Hubble expansion is the so-called dark energy, (70%). It is homogeneously distributed as opposed to cold dark matter, which is associated with the galaxies. Moreover its density remains relatively constant in a range of  $z$  values of  $O(1)$ . For these reasons most astrophysicists and cosmologists consider dark energy to have the characteristics of a cosmological constant, probably associated with some form of vacuum energy.

We are then presented with the seemingly enigmatic situation that, whereas one appeals to a phenomenological  $\Lambda$  to explain cosmogenesis and inflation, of planckian origin and magnitude, [1, 2], at present we are confronted with  $\Lambda$  of a vastly different order of magnitude ( $< 10^{-100}$ ). Do these phenomenologies have anything to do with each other? In this paper we argue in favor of the hypothesis that they are in fact strongly related. The present value of  $\Lambda$  could well be a remnant of inflation that is induced by fluctuations of vacuum energy. After the macroscopic fluctuation, that is responsible for inflation, decays into quanta, [2], there remain fluctuations of vacuum energy in the presence of these quanta (present day particles). We reinterpret the causet scenario of [3] in this light.

Our paper is laid out as follows: We first summarize the causet mechanism [3] in preparation for its reinterpretation and then make contact with inflation and our previous concept of the inflaton. We will also make contact with the creation of cisplanckian modes, which one supposes must occur to keep constant, in the mean, the cisplanckian mode density (equivalent to a constant planckian cut-off), see [4]. The analogous scenario of mode creation in the black hole evaporation process is discussed as well.

When the inflationary epoch is terminated, our interpretation of the causet mechanism does not modify the quantitative realization of [3] at least at this stage of development. This latter is given in terms of the parameter,  $\alpha$  (see Eq.1 below), the value of which is rather tightly constrained on phenomenological grounds [3]. This pa-

rameter plays a role in the estimate of the inflaton mass, using notions taken from [5].

It is evident, that like all phenomenologies which involve quantum gravity and planckian physics, this paper is conjectural in character. Its interest is to show that it is possible to formulate hypotheses that unify cosmogenesis, inflation, dark energy and mode creation.

A brief resume of [3] is now sketched. Random fluctuations of planckian units of energy (better, of action) are sequentially fed into the expanding cosmos. In ref. 3 these lumps are taken to be planckian elements of space-time as well. They are distributed at random with a mean planckian density. A stochastic process is set up in time in which the events in the  $i$ 'th slice of time generates those in the  $(i + 1)$ th [Eq.1 below]. This set of events is called a causet. Its observable effects are limited to the past light cone of a comoving observer hence proportional to the space time volume ( $V$ ) of the light cone bounded sufficiently far back in the distant past so as to validate the estimate  $V \propto H^{-4}$  where  $H(t)$  is the Hubble constant at the proper time of the observer. Two essential assumptions are:

1. Designating  $\rho_M$  as the energy density of everything else but the vacuum energy density ( $\rho_\Lambda$ ), it is assumed that as  $\rho_M \rightarrow 0$  then  $\rho_\Lambda \rightarrow 0$ . This is tantamount to the assumption that there is a stable quiescent vacuum towards which the universe evolves at late times; complete rest is the "natural" condition of the cosmos. This assumption is natural in that, were the vacuum state  $\rho_M = 0$  characterized by nonvanishing  $\rho_\Lambda$ , the universe in which this vacuum exists would expand, thereby creating energy,  $\rho_M$ , which is nonvanishing due to the temporal dependence of the metric.

For this very reason it is necessary to modify the first assumption somewhat. The final target state is metastable. We assume, as in [2], that large scale fluctuations, those necessary to be seized upon by the cosmological component of gravity to inflate and seed a new universe, are very rare.

2. Since  $\Lambda$  varies in time (and in a more detailed formulation it would vary in both space and time), there

would be sources of energy-momentum in general relativity which compensate for this variation. These would appear in cosmology in the equation for the acceleration of  $a$  ( $a$  = scale factor). This equation is dropped in [3] and will be dropped in the foregoing as well.

On the contrary the energy constraint (Friedmann equation) is maintained, an equation of energy balance between energy carried by gravity due to the expansion and energy sources due to other fields both in vacuum and in quanta. This equation is a result of invariance under time reparametrization, a physical principle one would loathe to give up.

We remark here on the general point of view given in [3] and which we shall adopt in the present work. The system treated is open. In [3] the causet elements arise from “nowhere” but are created causally (sequentially) from those already present. “Nowhere” can also be taken to mean from an unknown somewhere and this is the substance of our reinterpretation. The parameter  $\alpha$  is then interpretable as the squared matrix element of the transition from this “somewhere” to the cosmos accompanied by energy exchange between the two. This is what allows us to express the inflaton mass in terms of  $\alpha$ .

Our interpretation of [3] is perhaps more modest, seemingly less audacious than the creation of space time lumps in its quest for quantum gravity. We suppose that space-time is preexistent. The planckian lumps are the localized scenes where energy exchange occurs between the transplanckian world (i.e. degrees of freedom whose length scale is less than planckian, often called space time foam) and the cisplanckian world, our cosmos, the quanta therein and the vacuum in which these quanta are situated. This point of view allows one to conceive of inflation, mode creation and the present day  $\Lambda$  on a common basis.

It is commonly believed that the modes used to quantize fields in usual (cisplanckian) quantum field theory supply a valid description on scales of momenta less than planckian. ( $p < 1$ ) where we use planckian units. Herein there is an inherent frame dependence taken to be the cosmological rest frame, a weakness one hopes to eliminate in a future covariant formulation if such is possible. For  $p > 1$ , gravitational interactions among the modes become the dominant component of energy. One expects these modes to become overdamped and field theory on this scale must be considered a strong coupling theory. A possibility that is often envisaged is that fields on these short length scales fold up into localized structures of planckian and transplanckian dimensions and densities, Wheeler’s so-called foam.

The foam, in this image, serves as a reservoir of cisplanckian modes which we now explain. As the universe expands, wavelengths expand with it. Therefore a cut off of modes at  $p = O(1)$  at earlier times, say  $t = t_0$ , becomes reduced by a factor  $(a(t_0)/a(t))$ .

From inflationary times to the present this can result

in a reduction in mode density  $O(10^{-150})$ . But it is natural to suppose that cisplanckian physics does not vary sensibly in time in which case modes must be produced constantly. One possibility is that space is produced from “nowhere” as in causet theory (see [3]) and, in that space, modes of density  $O(1)$  exist because of the discretization. Another possibility is that space is preexistent. This is the hypothesis that has been explored in [5] to explain the phenomenological inflaton and its mass at the same time as mode creation. In this scenario the degrees of freedom that are locked into the localized transplanckian structures are steadily solicited so as to maintain the cisplanckian mode density constant as the universe expands. Vacuum is conceived as a sort of state of phase equilibrium between two fluids, the transplanckian reservoir and the cisplanckian world. There must always be as many degrees of freedom in the reservoir as is necessary to maintain this situation. An example would be supplied by the massy and/or short wave length modes of string theory if ever that hypothesis could be made to work. In fact, [6] supplies an image of high energy strings connected to black hole structures which could serve as a nice model of foam along these lines.

This cis-trans equilibrium thus is implemented by the exchange of degrees of freedom from planckian lumps of energy to and from modes. Localized structures of foam give and take energy from the mean situation wherein the cisplanckian vacuum energy density is maintained constant in the mean. As in any equilibrium, there will be fluctuations about this mean. These will carry both signs of energy and a sensible stochastic hypothesis for these fluctuations could be Eq.1 wherein the fluctuation of action at time  $t = t_{i+1}$  is preconditioned by its value at time  $t = t_i$ . Independently of any detailed mechanisms, as pointed out in ref 3, the order of magnitude of vacuum energy density at time  $t$ , in a universe where matter is present, is  $\pm[H(t)]^2$  since the volume of the backward light cone is  $O[(H(t))^{-4}]$  and the net energy transfer in this light cone is  $\pm N(t)^{1/2}$  where  $N(t)$  is the number of cis-trans exchanges which are supposed to be planckian in energy as well as density. Thus,  $N(t)$  is  $O[(H(t))^{-4}]$  as well, whence  $\rho_\Lambda \propto O(N^{1/2}/H^{-4}) = H^2(t)$ . As stated, our reinterpretation maintains this remarkably simple and successful explanation of  $\rho_\Lambda$ .

There is one very important point that must be emphasized, to wit: it is supposed that only the cisplanckian energy fluctuations result in the metric back reaction of general relativity here given by the variation for the Hubble constant appearing in the energy constraint. Degrees of freedom locked into the transplanckian structures do not have the long wavelength components available to affect a macroscopic back reaction. Indeed it is not even clear that Einsteinian general relativity as we usually practice it applies at all to the transplanckian world. One guesses it would not.

We may understand the above statement in a transpar-

ent way by considering theories of black hole evaporation that tame the transplanckian problem through mode production or less dramatically through conversion of over damped to under damped degrees of freedom ([7, 8]).

When one tracks an evaporated Hawking photon backwards in time one finds that its trajectory hugs the horizon at exponentially short lengths hence describing a vacuum fluctuation in the Schwarzschild metric which has exponentially large proper energy (eigen value of  $\delta/\delta r$ ,  $r$  being the proper radial coordinate). Thus the description in terms of modes fails in this region. Rather one must suppose that there is a planckian skin containing transplanckian structures about the horizon which feeds into the mode description once the fluctuation of the outgoing configuration ( in a wave packet description) is outside the skin. Then by subsequent Bogoljubov transformation, this fluctuation in part becomes an outgoing quantum and the energy constraint results in a back reaction of the metric described by a loss of black hole mass given by the frequency of that quantum in the asymptotic region. The transplanckian reservoir serves only to feed the cisplanckian modes and the metric back reaction is the back reaction to this latter part only. We suppose that something of this sort also happens in the cosmological expansion wherein the cisplanckian field configuration arises out of the transplanckian structure due to expansion.

On the basis of this example, one may develop a predilection in favour of our interpretation that causets do not concern the creation of space in lumps, but only the exchange of energy associated with these lumps. Indeed, in the black hole example, space is always present or presumed as such in all existing models. The transplanckian skin about the horizon in the type of model we are discussing is the locus of sites of cis trans exchange. Note that if it turns out that the correct black hole transplanckian taming is in-out scattering, as advocated for example by G. 't Hooft, then this example will be irrelevant.

Let us now turn to inflation. The parameter  $\alpha$  in [3] is defined through the recursion relation Eq.5 in [3].

$$\rho_{\Lambda,i+1} = \frac{S_{i+1}}{N_{i+1}} = \frac{S_i + \alpha \xi_{i+1} (\delta N_i)^{1/2}}{N_i + \delta N_i} \quad (1)$$

The index  $i$  denotes the  $i$ 'th slice through the backward light cone at time  $t_i$ ,  $N_i$  is the total number of lumps accumulated at time  $t_i$  in that cone.  $\xi_i$  is a random variable of mean zero and standard deviation 1.  $\delta N_i$  is the number of lumps in the  $i$ th slice.  $\alpha$  is a numerical parameter. Good agreement with phenomenology was found in [3] with  $\alpha = O(10^{-2})$ . See [3] for further detail and the understanding why  $\alpha$  must be constrained to  $O(10^{-2})$ . It is no small achievement that there exist runs which seem to be accurate replicas of phenomenology back to  $z$  values of  $O(1)$ .

We have presented these details to motivate our interpretation of  $\alpha$  as the squared matrix element for cis trans exchange of a planckian unit of energy between the transplanckian lump in the reservoir and the cisplanckian world.

The relevance for inflation and the mass of the inflaton is now presented. As in Refs.[1, 2], we start at some time when space is flat hence  $\rho_M = 0$ , and  $\rho_\Lambda$ , in the mean, is 0 as well. Consistent with the assumption of our target values as discussed above. this vacuum state is taken to be an equilibrium between the cis and trans fluids.

Though small scale length fluctuations regress, there exist large scale fluctuations which will inflate. (See [2]). It was argued in [2] that this large fluctuation is the phenomenological inflaton often represented as a scalar field but which is commonly believed to be a manifestation of quantum gravity. In the usual inflationary scenarios this macroscopic fluctuation will slowly regress after the number of e-folds is sufficient to account for our observed universe. The initial fluctuation is chosen sufficiently large to do the job [2]. Of course, small scale fluctuations accompany this regression. In current phenomenology one accounts for the spectrum of cosmic background radiation in terms of one class of these fluctuations. But we now are appealing to another class, those that arise explicitly from the fluctuations of the cis trans exchange equilibrium and which is our interpretation of [3]. The big difference between the physics of the macroscopic fluctuation responsible for inflation and the microscopic fluctuations which have been discussed until now is that these latter occur in a universe already filled with energy. As a result, the cisplanckian system can both take on and give up energy in these exchanges. This is the natural and beautiful explanation of [3] for why  $\rho_\Lambda = O(H^2)$  at present. On the contrary, the macroscopic fluctuation called the inflaton is a phenomenon which takes place when  $\rho_M = 0$ . The cisplanckian system apart from microscopic fluctuations then has but negligible energy to give up during the evolution of the large macroscopic fluctuation. Almost all of the transplanckian lumps within this large fluctuation only give energy to the cisplanckian degrees of freedom i.e. the modes, thereby creating our universe after reheating. As we have emphasized the macroscopic metric back reaction encoded in the expanding scale factor is due only to changes in energy in the cisplanckian world.

We thus understand why the inflationary Hubble constant, or equivalently the value of  $\rho_\Lambda$  during inflation is of planckian scale. All the transplanckian lumps that give up energy act in consort to increase the total energy of the inflating universe until towards the end when the inflatonic motor runs out; preheating takes place and the isentropic expansion ensues. Thus the energy density induced in the cisplanckian world at inflationary times is no longer  $O(N^{1/2}/V)$  [where we recall that  $N$  is the number of lumps that affect the trans cis exchange and  $V$  is the

space time volume of the backward light cone]. Rather in the inflationary epoch the energy density passed to the cisplanckian world is  $O(N/V)$  [ $= (H^{-4}/H^{-4}) = O(1)$ ]. In short, inflation is a process of coherent passage to the cisplanckian world of energy from transplanckian lumps whereas the dark energy is a random incoherent process with either sign of energy exchange. At the same time, and presumably by the same mechanism cisplanckian modes are created from the transplanckian degrees of freedom.

One may refine somewhat the estimate of the order of magnitude of  $\rho_\Lambda$  during inflation or at least the effective inflaton mass by appeal to  $\alpha$  and its interpretation as the squared matrix element for the exchange.

It was argued in [5] that the inflaton is a wave of varying density induced by fluctuations from the equilibrium resulting from cis trans exchanges. On this basis the inflaton mass was estimated from the equation

$$\mu^2 = pM^2 \quad (2)$$

where  $p$  = probability of cistrans exchange and  $M$  equal mass scale of the matrix element for the exchange. We here identify  $p = \alpha$  ( $= O(10^{-2})$ ). Phenomenologically,  $\mu = O(10^{-6} - 10^{-5})$  whence  $M = O(10^{-4})$ , a not unreasonable number but nevertheless rather lower than what one might have expected. Only when a fundamental theory of planckian and transplanckian physics begins to take on a credible form will we be in a position to confirm or infirm these considerations.

After the first draft of this article I had occasion to hear Renaud Parentani present his recent work on the production of cis planckian modes from a transplanckian reservoir field. The essential input of his theory is a set of stochastic assumptions governing the reservoir field,

designed to give the desired result of a mean planckian density of cis planckian modes during the Hubble expansion. The output is the elegant passage from over to under damping of degrees of freedom at the planckian level. It is clear that this point of view has much in common with ideas expressed in the above paragraphs and it will be most interesting to see if this recent work can furnish more quantitative detail on our point of view of dark energy and inflation.

**Acknowledgement:** It is a pleasure for me to acknowledge with gratitude the friendly patient explanations of Raphael Sorkin, of the beautiful ideas expressed in [3]. During this period I was hosted by the Perimeter Institute, Waterloo, Ontario, Canada. I wish to express my gratitude to the many people who helped me at Perimeter and my appreciation of the productive and friendly atmosphere that prevails there; in particular, Achim Kempf has been most helpful and kind.

- 
- [1] R. Brout, F. Englert, E. Gunzig, *Gen. Rel. Grav.* **10** 1 (1979) (1st Prize Gravity Award Essay), A.A. Starobinski, *Phys. Lett.* **91B** 99 (1980)
  - [2] E.W. Kolb, M.S. Turner, *The Early Universe*, Perseus (1993)
  - [3] M. Ahmed, S. Dodelson, P.B. Greene, R. Sorkin, *Phys. Rev.* **D69** 103523 (2004)
  - [4] A. Kempf, *Phys. Rev.* **D63** 083514 (2001), B. Z. Foster, T. Jacobson, *JHEP* 0408:024 (2004)
  - [5] R. Brout, gr-qc/0201060
  - [6] T. Damour, G. Veneziano, *Nucl. Phys.* **B568** 93 (2000)
  - [7] R. Brout, C. Gabriel, M. Lubo, P. Spindel, *Phys. Rev.* **D59** 044005 (1999)
  - [8] C. Barrabes, V. P. Frolov, R. Parentani, *Phys. Rev.* **D59** 124010 (1999), *Phys. Rev.* **D62** 044020 (2000)